arXiv:1407.4771v1 [math.GR] 17 Jul 2014

ON VERTEX-UNIPRIMITIVE NON-CAYLEY GRAPHS OF ORDER PQ

MOHAMMAD A. IRANMANESH

ABSTRACT. Let p and q be distinct odd primes. Let $\Gamma = (V(\Gamma), E(\Gamma))$ be a non-Cayley vertex-transitive graph of order pq. Let $G \leq \mathrm{Aut}(\Gamma)$ acts primitively on the vertex set $V(\Gamma)$. In this paper, we show that G is uniprimitive which is primitive but not 2-transitive and we obtain some information about p, q and the minimality of the Socle $T = \mathrm{Soc}(G)$.

1. INTRODUCTION

Let $\Gamma = (V(\Gamma), E(\Gamma))$ or simply Γ be a finite, simple and undirected graph. We denote by $\mathrm{Aut}(\Gamma)$, the automorphism group of Γ and we say that Γ is vertex-transitive if $\mathrm{Aut}(\Gamma)$ acts transitively on $V(\Gamma)$. The cardinality of $V(\Gamma)$ is called the order of the graph. For a group G and a subset S of G such that $1_G \notin S$ and $S^{-1} = S$, the Cayley graph $\Gamma = \mathrm{Cay}(G, S)$ is defined to have vertex set G and for $g, h \in G, \{g, h\}$ is an edge if and only if $gs = h$ foe some $s \in S$. Every Cayley graph is a vertex-transitive graph and conversely, see [1], a vertex-transitive graph is isomorphic to a Cayley graph for some group if and only if its automorphism group has a regular subgroup on vertices.

By $\mathcal{NC}$ we denote the set of positive integers n for which there exists a vertex-transitive non-Cayley graph on n vertices which is not a Cayley graph. In [9], Marušič proposed the determination of $\mathcal{NC}$ and in [10], he proved that $p, p^2, p^3 \notin \mathcal{NC}$ for prime p. Mckay and Praeger settled the status of all non-square-free numbers n are in $\mathcal{NC}$, with the exceptions $n = 12, n = p^2, n = p^3$, p prime and they finished the characterization of non-Cayley numbers which are the product of two distinct primes (see [7, 8]). The following theorem characterize all numbers $n = pq \in \mathcal{NC}$ where p, q are distinct primes.

1991 *Mathematics Subject Classification.* 20B20, 05C25.
Key words and phrases. Uniprimitive group, Non-Cayley graph, Socle.

Theorem 1. [8, Theorem 1] *Let $p < q$ be primes. Then $pq \in \mathcal{NC}$ if and only if one of the following holds.*

(1) *p^2 divides $q-1$;*
(2) *$q = 2p-1$ or $q = \frac{p^2+1}{2}$;*
(3) *$q = 2^t+1$ and either p divides 2^t-1 or $p = 2^{t-1}-1$;*
(4) *$q = 2^t-1$ and $p = 2^{t-1}+1$;*
(5) *$pq = 7.11$.*

For characterization of non-Cayley number which are the product of three distinct primes we refer to [13, 4, 5].

Suppose that a group G acting on a set V transitively. We say that a partition Σ of V is G-invariant if G permutes the blocks of Σ. A transitive action of G on V is said to be primitive if all G-invariant partitions of V are trivial. A primitive permutation group which is not 2-transitive is said to be uniprimitive. Also a transitive permutation group is said to be minimal transitive if all of its proper subgroups are intransitive. Suppose that Γ is a graph of order pq. In this paper we prove that if $G \leq \mathrm{Aut}(\Gamma)$ acts primitively on the vertex set of Γ, then either Γ is a Cayley graph or G is uniprimitive and when $pq \notin \mathcal{NC}$ then $T = \mathrm{Soc}(G)$ is not minimal transitive.

2. Primitive permutation groups of degree pq

First, we investigate primitive permutation groups of order pq which are 2-transitive.

Proposition 2. *Let p and q be distinct odd primes such that $p < q$ and suppose that G, a subgroup of S_{pq}, is 2-transitive of degree pq with Socle T. Then T is a non-abelian simple group and is one of the group listed below.*

(*i*) *$T = A_{pq}$;*
(*ii*) *$T = \mathrm{PSL}(n,s)$ and $pq = \frac{s^n-1}{s-1}$;*
(*iii*) *$T = \mathrm{PSU}(3,2^a)$ and $p = 2^a+1, q = 2^{2a}-2^a+1$;*
(*iv*) *$T = \mathrm{Sz}(8)$ and $p = 5, q = 13$;*
(*v*) *$T = A_7$ and $p = 3, q = 5$.*

Also in case (i) and (ii) for some but not all p and q, we have $pq \in \mathcal{NC}$. In case (iii), $pq \notin \mathcal{NC}$ except where $p = 5, q = 13$ and in case (v), $pq \notin \mathcal{NC}$ and in case (iv), $pq \in \mathcal{NC}$.

Proof. By the "O'Nan-Scott Theorem" [12], G is almost simple, that is, G has a unique minimal normal subgroup T which is a non-abelian simple group. All possibilities for the socle of almost 2-transitive groups are given for example in [2]. Note that in case (iv) we have the Suzuki groups $\mathrm{Sz}(q)$ with $s = 2^{2a+1} = \frac{r^2}{2} \geq 8$. In this case $pq = s^2 + 1 = (s + r + 1)(s + r - 1)$. Thus 5 divides pq and hence $p = 5$ and $q = 1$. From Theorem 1 we conclude that in case (iv), $pq \in \mathcal{NC}$ and in case (iii), $pq \in \mathcal{NC}$ if and only if $a = 2, p = 5$ and $q = 13$. □

Now we deal with the primitive permutation groups of degree pq which are not 2-transitive, that is, the uniprimitive permutation groups of degree pq.

Proposition 3. *Let p and q be distinct odd primes such that $p < q$. Suppose that G, is a uniprimitive permutation subgroup of S_{pq}, of degree pq with Socle T. Then T is a non-abelian simple group and Table 1 contains all the possibilities for T, p and q.*

Proof. A similar argument to the proof of Proposition 2, we conclude that T is a non-abelian simple group. All possibilities for the socle of uniprimitive permutation groups of degree pq are given in [11] based on [6] which are listed in Table 1. □

T	q	p	T	q	p
A_q	$q \geq 5$	$\frac{q-1}{2}$	$\mathrm{PSL}(2, 19)$	19	3
A_{q+1}	$q \geq 5$	$\frac{q+1}{2}$	$\mathrm{PSL}(2, 29)$	29	7
A_7	7	5	$\mathrm{PSL}(2, 59)$	59	29
$\mathrm{PSL}(4, 2)$	7	5	$\mathrm{PSL}(2, 61)$	61	31
$\mathrm{PSL}(5, 2)$	31	5	$\mathrm{PSL}(2, 23)$	23	11
$\mathrm{PSp}(4, 2^a)$	$2^{2a} + 1$	$2^a + 1$	$\mathrm{PSL}(2, 11)$	11	5
$\Omega^{\pm}(2n, 2)$	$2^n \mp 1 > 7$	$2^{n-1} \pm 1$	$\mathrm{PSL}(2, 13)$	13	7
$\mathrm{PSL}(2, q)$	$q \geq 13$	$\frac{q+1}{2}$	M_{23}	23	11
$\mathrm{PSL}(2, q)$	$q \geq 7$	$\frac{q-1}{2}$	M_{22}	11	7
$\mathrm{PSL}(2, p^2)$	$q = \frac{p^2+1}{2}$	p	M_{11}	11	5

TABLE 1. Socle of uniprimitive groups of degree pq.

3. Uniprimitive graphs of order pq

In this section we prove that there is no any non-Cayley graph of order pq which admits a 2-transitive subgroup of automorphisms.

Theorem 4. *Let p and q be distinct odd primes and $\Gamma = (V(\Gamma), E(\Gamma))$ be a vertex-transitive graph of order pq. Suppose that there exists a subgroup G of $\mathrm{Aut}(\Gamma)$ which acts 2-transitively on $V(\Gamma)$. Then Γ is a Cayley graph.*

Proof. Let Γ be a vertex-transitive graph of order pq and suppose that $G \leq \mathrm{Aut}(\Gamma)$ is 2-transitive on $V(\Gamma)$. If Γ be an empty graph of order pq, then obviously Γ is a Cayley graph. So we may assume that there exist $x, y \in V(\Gamma)$, such that $\{x, y\} \in E(\Gamma)$. Since G is doubly transitive on $V(\Gamma)$, there exists $g \in G$, such that $x^g = x'$ and $y^g = y'$. Thus $\{x, y\}^g = \{x', y'\} \in E(\Gamma)$. This shows that Γ is a complete graph and the proof is complete. □

Corollary 5. *Let Γ be a non-Cayley vertex-transitive graph of order pq. If $G \leq \mathrm{Aut}(\Gamma)$ acts primitively on $V(\Gamma)$, then G is uniprimitive and $T = \mathrm{Soc}(G)$ is one of the group listed in Table 2. Also if $pq \notin \mathcal{NC}$, then T is not minimal transitive.*

Proof. The first part is direct consequence of Theorem 4. If $pq \notin \mathcal{NC}$, then T is one of the groups in Table 2. For each line we see that T has a transitive subgroup isomorphic to a Frobenius group of order pq. This can be shown by using [3]. □

T	q	p
A_q	$q \geq 5$	$\frac{q-1}{2}$
$\mathrm{PSL}(5,2)$	31	5
$\mathrm{PSL}(2,q)$	$q \geq 7$	$\frac{q-1}{2}$
$\mathrm{PSL}(2,29)$	29	7
$\mathrm{PSL}(2,59)$	59	29
$\mathrm{PSL}(2,23)$	23	11
$\mathrm{PSL}(2,11)$	11	5
M_{23}	23	11
M_{11}	11	5

Table 2. Socle of uniprimitive groups of degree pq where $pq \notin \mathcal{NC}$.

Acknowledgment. The author wishes to thank Yazd University Research Council for financial support.

References

[1] N. L. Biggs, *Algebraic grapph theory*, Cabridge University Press, New York, (1974).

[2] P. J. Cameron, Finite permutation groups and finite simple groups, *Bull. London Math. Soc.*, **13**, (1981), 1−22.

[3] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker and R. A. Wilson, *An Atlas of finite groups*, Clarendon Press, Oxford, (1985).

[4] A. Hassani, Mohammad A. Iranmanesh, C. E. Praeger, On vertex-imprimitive graphs of order a product of three distinct odd primes. *J. Combin. Math. and Combin. Computing*, **28**, (1998), 187−213.

[5] Mohammad A. Iranmanesh, C. E. Praeger, On non-Cayley vertex-transitive graphs of order a product of three primes. *J. Combinatorial Theory, Series B*, **81**, (2001), 1−19.

[6] M. W. Liebeck, J. Saxl, Primitive permutation groups containing an element of large prime order. *J. London Math. Soc.*, (2) **31**, (1985), 237−249.

[7] B. D. Mckay, C. E. Praeger, Vertex-transitive graphs which are not Cayley graphs I, *J. Austral. Math. Soc.*, **56**, (1994), 53−63.

[8] B. D. Mckay, C. E. Praeger, Vertex-transitive graphs which are not Cayley graphs II, *J. Graph Theory*, **22**, (1996), 321−334.

[9] D. Marušič, Cayley properties of vertex symmetric graphs, *Ars. Combin.*, **16B**, (1983), 297−302.

[10] D. Marušič, Vertex-transitive graphs and digraphs of order p^k, *Ann. Disc. Math.*, **27**, (1985), 115−128.

[11] D. Marušič, R. Scapellato and B. Zgrablič, On quasiprimitive pqr-graphs, *Algebra Coll.*, **2**, (1995), 295−314.

[12] C. E. Praeger, An O'Nan-Scott Theorem for finite quasiprimitive permutation groups, and an application to 2-transitive groups, *J. London. Math. Soc.*, (2) **47**, (1993), 227−239.

[13] A. Seress, On vertex-transitive non-Cayley graphs of order pqr, *Discrete Math.*, **182** (1998), 279−292.

M. A. Iranmanesh, Department of Mathematics

Yazd University, Yazd, 89195-741 , Iran

E-mail address: iranmanesh@yazd.ac.ir